

Saturation Effects in the VLF Scattering off HF Heated Ionosphere

Nikolai G. Lehtinen, Timothy F. Bell and Umran S. Inan

VLF Group, Stanford University
Stanford, CA

National Radio Science Meeting
Boulder, CO

January 6, 2012



Outline

- 1 Overview
- 2 Stanford Full-Wave Method (SFWM) code
- 3 Scattering with the method of moments (MoM)
 - Model description
 - Results



Outline


- 1 Overview
- 2 Stanford Full-Wave Method (SFWM) code
- 3 Scattering with the method of moments (MoM)
 - Model description
 - Results

VLF scattering on *D*-region disturbances

The VLF perturbations are caused by *D*-region disturbances due to HF heaters and can be calculated using Earth-ionosphere waveguide mode theory:

- with **WKB** and **Born** approximations [Barr *et al*, 1985; Demirkol, Ph. D thesis, 1999].
- with **Born** but no **WKB** [Lehtinen *et al*, 2011]
- neither **Born** nor **WKB** [Foust *et al*, 2011; present work]

We use Stanford Full-Wave Method (SFWM) together with the method of moments (MoM), which uses less computer resources than discontinuous Galerkin (DG) finite element method [Foust *et al*, 2011].




VLF scattering by an HF heater: NLK/HAARP

NLK VLF transmitter:

- Modelled as a ground-based vertical dipole
- $f = 24.8$ kHz
- $P = 250$ kW

HAARP HF heater:


- $f_{\text{HF}} = 5$ MHz
- ERP = 1 GW
- Beam width ~ 23 km [*Payne et al, 2007*], we assume Gaussian horizontal shape
- ΔT_e and Δv_e are found using kinetic equations



Incident VLF wave


Strongest modes at $R_0 = 2000$ km (disturbance)

- Modes are calculated using SFWM using night-time ionosphere
- Attenuation is due to both **absorption** and **radiation into ionosphere**



Change in ν_e due to heating

Kinetic model results for steady heating starting at $t = 0$





Outline

- 1 Overview
- 2 **Stanford Full-Wave Method (SFWM) code**
- 3 Scattering with the method of moments (MoM)
 - Model description
 - Results


Stanford Full-Wave Method (SFWM) code

Capabilities:

- Arbitrary **plane stratified** medium, e.g., a horizontally-stratified magnetized plasma with an arbitrary direction of geomagnetic field (such as ionosphere)
- Arbitrary configuration of harmonically varying currents
- Provides full wave 3D solution of both whistler waves launched into ionosphere and VLF waves launched into Earth-ionosphere waveguide
- Stable against the “swamping” instability by evanescent waves
- Efficient use of the computer resources, easily parallelized

Applications:

- Trans-ionospheric propagation
- Earth-ionosphere waveguide propagation
- Scattering on *D*-region disturbances



We work in Fourier (horizontal wave vector \mathbf{k}_\perp) domain:

- 1 For each $\mathbf{k}_\perp = \text{const}$ (Snell's law) \implies find k_z , E and H in each layer for each of 4 plane wave modes (2 up, 2 down)
- 2 Use continuity of \mathbf{E}_\perp and \mathbf{H}_\perp between layers to find reflection coefficients $R^{u,d}$ and mode amplitudes \mathbf{u} , \mathbf{d}
 - Recursion order $R_{k+1}^u \rightarrow R_k^u$ and $\mathbf{u}_k \rightarrow \mathbf{u}_{k+1}$ provides stability against “swamping” of solution by evanescent waves
 - Represent source currents as boundary conditions on \mathbf{E}_\perp and \mathbf{H}_\perp between layers
- 3 Inverse Fourier transform from \mathbf{k}_\perp to \mathbf{r}_\perp

Outline

- 1 Overview
- 2 Stanford Full-Wave Method (SFWM) code
- 3 Scattering with the method of moments (MoM)
 - Model description
 - Results



Outline


- 1 Overview
- 2 Stanford Full-Wave Method (SFWM) code
- 3 Scattering with the method of moments (MoM)
 - Model description
 - Results



Previously used Born approximation

- Neglect the scattered field \mathbf{E}_s compared to the incident field \mathbf{E}_0 inside the perturbed region
- \mathbf{E}_0 acting together with the perturbation $\Delta\hat{\sigma}$ creates currents which radiate \mathbf{E}_s
- What if \mathbf{E}_s is comparable to \mathbf{E}_0 ?

Motivation: $\Delta\hat{\sigma}$ may be large



Description of the method of moments (MoM)

- Green's function is a 3×3 matrix \hat{G} with components

$$G_{ij}(\mathbf{r}_o, \mathbf{r}_s) \equiv E_i(\mathbf{r}_o) \text{ created by current } \mathbf{J}(\mathbf{r}) = \hat{x}_j \delta(\mathbf{r} - \mathbf{r}_s)$$

\mathbf{r}_s — source position

\mathbf{r}_o — observer position

In our case, Green's function is in the stratified medium, and currents $\mathbf{J} = \Delta \hat{\sigma} \mathbf{E}$ are due to conductivity perturbation.

- We have an integral equation for the scattered field \mathbf{E}_s :

$$\mathbf{E}_s(\mathbf{r}) = \int \hat{G}(\mathbf{r}, \mathbf{r}') \Delta \hat{\sigma}(\mathbf{r}') [\mathbf{E}_0(\mathbf{r}') + \mathbf{E}_s(\mathbf{r}')] d^3 \mathbf{r}'$$

where the integration is over the perturbed region ($\Delta \hat{\sigma} \neq 0$).

- MoM** makes use of discretisation of \mathbf{J} , \mathbf{E} . Then the integral equation is solved numerically, and involves an inversion of a large matrix.




Outline


- 1 Overview
- 2 Stanford Full-Wave Method (SFWM) code
- 3 Scattering with the method of moments (MoM)
 - Model description
 - Results

3D calculation of E_z from scattering of QTM1 mode


Vertical slice





Horizontal slice




Error in $\Delta\mathbf{J} = \Delta\hat{\sigma}\mathbf{E}$ due to Born approximation



Amplitude change on the ground



Upward flux change at 137.5 km





Conclusions:

- The previously calculated scattering in Born approximation overestimates the effect of strongly heated ionosphere;
- Quantitatively the results are still of the same order.

Acknowledgements:

This work was supported by Dept. of Air Force grant FA9453-11-C0011 and DTRA grant HDTRA1-10-1-0115 to Stanford University.